



Osmocote[®]
Exact
Tablet

Tablet 8-9M

Long-lasting nutrition, delivered with precision

14 | 3.5 | 9.1 | TE
N | P | K



Guaranteed analysis

Elemental

N	Total Nitrogen	14%
	Nitrate nitrogen (N-NO ₃)	6.2%
	Ammoniacal nitrogen (N-NH ₄)	7.8%
P	Phosphorus	3.5%
	Water soluble (P)	2.6%
K	Potassium	9.1%
	Water soluble (K)	9.1%
Mg	Magnesium	1.2%
	Water soluble (Mg)	0.8%
B	Boron	0.01%
	Water soluble (B)	0.01%
Cu	Copper	0.046%
	Water soluble (Cu)	0.028%
Fe	Iron	0.41%
	Iron EDTA (Fe)	0.08%
Mn	Manganese	0.05%
Mo	Molybdenum	0.018%
	Water soluble (Mo)	0.012%
Zn	Zinc	0.013%

Description

Osmocote[®] Exact Tablets 8-9M deliver accurate doses of nutrients with remarkable ease. Their cone-like shape means the tablet can be pushed down easily into growing medium. The water-soluble adhesive dissolves and the tablets break apart. The tablets have a longevity of 8 to 9 months at 21°C and come in 5 g or 7.5 g.

Benefits

- Simple and safe application: each pot receives the same dosage
- High nutrient efficiency thanks to accurate dosage
- Tablets stay locked in the growing medium, so no loss of fertilizer

How to use

- 1 For application in existing cultures, push tablet down completely into the growing medium.
- 2 In hanging baskets, arrange the tablets in a circle in the growing medium around the plant.
- 3 If you're using drip irrigation, push the tablet in the growing medium directly under the dripper.
- 4 The temperature affects Osmocote® Exact Tablets' longevity, determined at 21°C. 16°C: 10-11M
21°C: 8-9M 26°C: 6-7M.
- 5 Store under dry conditions.
- 6 Properly seal partly used or damaged bags.

Application rates

	Light feeding	Normal feeding	When 50% of nutrition is supplied by Peters or Universol
Container Nursery Stock, Pot and Bedding plants	5 g/l	6 g/l	3 g/l

Important: Rates for Osmocote are based on pot volumes. When incorporating fertiliser throughout the media and repotting plants into bigger pots, the dosage rate should be increased to compensate for the dilution effect. Please contact your ICL advisor for plant-specific recommendations.

Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.