



Osmocote[®]
Exact
Standard

Standard 8-9M

Precision nutrition for 8-9 months

15 | 3.9 | 9.1 | TE
N | P | K



Guaranteed analysis

Elemental

N	Total Nitrogen	15%
	Nitrate nitrogen (N-NO ₃)	6.6%
	Ammoniacal nitrogen (N-NH ₄)	8.4%
P	Phosphorus	3.9%
	Water soluble (P)	2.8%
K	Potassium	9.1%
	Water soluble (K)	9.1%
Mg	Magnesium	1.2%
	Water soluble (Mg)	0.8%
B	Boron	0.023%
	Water soluble (B)	0.022%
Cu	Copper	0.060%
	Water soluble (Cu)	0.047%
Fe	Iron	0.45%
	Iron EDTA (Fe)	0.09%
Mn	Manganese	0.065%
Mo	Molybdenum	0.024%
	Water soluble (Mo)	0.024%
Zn	Zinc	0.028%
	Water soluble (Zn)	0.019%

Description

Safety and consistency: what more could you want?

Osmocote[®] Exact Standard 8-9M is the safest and most consistent way to feed your crop, tailored to your precise requirements. An easy way to make sure your plants get the nutrition they need throughout the season. The product has a percentage of coloured granules, to indicate their longevity too.' At tab 'Guaranteed analysis', can we go to elemental direct?

Benefits

- Perfect for crops that need a nutrient release period of 8-9M
- Safe: controlled release and high level of all essential trace elements
- Consistent: each bag gives identical results
- Tailored release pattern: exactly suited to your crop's requirements for optimal growth

How to use

- 1 The temperature affects the longevity of Osmocote® Exact Standard, determined at an overall average temperature of 21°C it will be 8-9M, at 16°C it will be 10-11M, at 26°C it will be 6-7M.
- 2 Close partly used or damaged bags securely.
- 3 Store under dry conditions.
- 4 If you need more information, please contact your technical support.

Application rates

	Light feeding	Normal feeding	When 50% of nutrition is supplied by Peters or Universol
Container Nursery Stock, Pot and Bedding plants	5 g/l	6 g/l	3 g/l

Important: Rates for Osmocote are based on pot volumes. When incorporating fertiliser throughout the media and repotting plants into bigger pots, the dosage rate should be increased to compensate for the dilution effect. Please contact your ICL advisor for plant-specific recommendations.

Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.