



## Lo.Start 16-18M

Feed your crop for more than a year

15 | 3.5 | 9.1 | TE  
N | P | K



## Guaranteed analysis

### Elemental

N	Total Nitrogen	15%
	Nitrate nitrogen (N-NO <sub>3</sub> )	6.6%
	Ammoniacal nitrogen (N-NH <sub>4</sub> )	8.4%
P	Phosphorus	3.5%
	Water soluble (P)	2.6%
K	Potassium	9.1%
	Water soluble (K)	9.1%
Mg	Magnesium	1.2%
	Water soluble (Mg)	0.8%
B	Boron	0.02%
	Water soluble (B)	0.02%
Cu	Copper	0.050%
	Water soluble (Cu)	0.031%
Fe	Iron	0.45%
	Iron EDTA (Fe)	0.09%
Mn	Manganese	0.06%
Mo	Molybdenum	0.020%
	Water soluble (Mo)	0.014%
Zn	Zinc	0.015%

## Description

Developed especially for slow-growing and sensitive crops, that stay in the same pot for more than one season.

Osmocote® Exact Lo.Start 16-18M gets your young plants off to a healthy start, releasing nutrition over more than a year. Ideal for situations where a slow start is required, such as slow rooting young plants and early potting in tunnels.

Osmocote Exact Lo.Start 16-18M is also suitable for warm climates. Your go-to precision nutrition solution to get slow-growers off to a flying start.

## Benefits

- \\ Nutrition supplied to young, slow-growing plants all year
- \\ Safe: controlled release and high level of all essential trace elements
- \\ Consistent: each bag produces identical results
- \\ Tailored release pattern suited to your crop's requirements

## How to use

- 1 The temperature affects Osmocote® Exact Lo.Start longevity, determined at 21°C. 16°C: 20-22M  
21°C: 16-18M 26°C: 12-14M.
- 2 Close partly used or damaged bags securely.
- 3 Store under dry conditions.
- 4 If you need more information, please contact your technical support.

## Application rates

	Light feeding	Normal feeding	When 50% of nutrition is supplied by Peters or Universol
Container Nursery Stock, Pot and Bedding plants	8 g/l	10 g/l	5 g/l

Important: Rates for Osmocote are based on pot volumes. When incorporating fertiliser throughout the media and repotting plants into bigger pots, the dosage rate should be increased to compensate for the dilution effect. Please contact your ICL advisor for plant-specific recommendations.

## Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.