



## Agroleaf® Liquid

### Iron

Iron out iron deficiency in your crops

0 | 0 | 0 | TE  
N P2O5 K2O



## Guaranteed analysis

Oxide		
N	Total Nitrogen	0%
P2O5	Phosphorus Pentoxide	0%
K2O	Potassium Oxide	0%
Fe	Iron	4.400%
	Water soluble (Fe)	4.400%
	Iron EDTA (Fe)	4.400%

## Description

Agroleaf® Liquid Iron is a premium foliar feed in liquid form that keeps iron deficiency at bay. It has been formulated for use in crops of all kinds, especially soft fruits. Agroleaf® Liquid Iron features ICL's unrivalled F3 SurfActive technology, which we designed specifically to make each foliar application more efficient. It does this in three ways: by improving spreading, adhesiveness, and retention. F3 technology reduces the surface tension of the droplets, allowing nutrients to distribute over the leaves more evenly for greater coverage and better nutrient uptake. It also retains more nutrients, preventing them from running or bouncing off waxy leaves. F3 aids nutrients in the formation of small deposits on the leaf surface, which are re-activated after rewetting. This prevents evaporation, which means the nutrients linger for longer, improving foliar application.

## Benefits

- \\ Keeps your plants nourished with the iron they need
- \\ Great for a wide range of crops, particularly soft fruits
- \\ Better spreading, adhesiveness, and retention

## How to use

1

If you need more information, please contact your technical support.

## Application rates

Crop recommendation	Dosage liter/ha	Water volume liter/ha	Conc (%)	Timing
Fruiting vegetables	3-6	300-600	1	During the entire crop cycle, 3-4 times
Strawberry	3-6	250-400	1.2-1.5	During the entire crop cycle, 3-4 times
Blueberry	3-6	250-400	1.2-1.5	During the entire crop cycle, 3-4 times
Other soft fruits	3-6 5-8	600-1000	0.5-0.6 0.8	During the entire crop cycle, 3-4 times for curative applications
Orchards	3-5 5-7	600-1000	0.5-0.6 0.8	During the entire crop cycle, 3-4 times for curative applications
Vineyards	3-5 5-7	600-1000	0.5-0.6 0.8	During the entire crop cycle, 3-4 times for curative applications
Ornamentals	2-3	600-1000	0.3	During the entire crop cycle, 3-4 times

\* Use lower dosages in lower volume of water/ha and higher dosages when the water volume is increased. Trial first on a small scale before changing the rate, or any other variables, As circumstances can differ and the application of our products is beyond our control, ICL cannot be held responsible for any adverse results.

## Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.