



**Osmocote®  
Topdress**

## Topdress 8-9M

Topdress in the fast lane

17 | 7 | 10 | TE  
N | P<sub>2</sub>O<sub>5</sub> | K<sub>2</sub>O



## Guaranteed analysis

Oxide		
N	Total Nitrogen	17%
	Nitrate nitrogen (N-NO <sub>3</sub> )	7.8%
	Ammoniacal nitrogen (N-NH <sub>4</sub> )	9.2%
P <sub>2</sub> O <sub>5</sub>	Phosphorus Pentoxide	7%
	Water soluble (P <sub>2</sub> O <sub>5</sub> )	6.1%
K <sub>2</sub> O	Potassium Oxide	10%
	Water Soluble (K <sub>2</sub> O)	10.0%
Fe	Iron	2.00%
	Water soluble (Fe)	2.00%

## Description

Osmocote® Topdress 8-9M both features Osmocote® coating for a slow release of nutrients over 8-9 months and gives your ornamentals a little helping of fast-release nitrogen and phosphorous straightaway. Osmocote® Topdress is packed with N, P, K, and iron. High levels of nitrogen and iron quickly bring extra green to your plants, and nurse them back to health if they've been lacking in nutrients. Specially developed for topdressing with container nursery stock. A state-of-the-art nutrition solution for optimum container stock performance.

## Benefits

- \\ Healthy green container stock fast
- \\ Instant nutrition and slow release in one
- \\ Dust-free

## How to use

- 1 Osmocote® Topdress has been specifically developed for topdress applications in container nursery stock.
- 2 You can use it to topdress container plants to quickly green them up or correct any deficiency symptoms.
- 3 Topdressing is easy if you use overhead irrigation or drip irrigation. If you use drip irrigation, place the product directly under the drippers.
- 4 You can apply Osmocote® Topdress with recommended ICL topdress applicators. Irrigate well after application. Ask your ICL advisor for more information.
- 5 Close partly used or damaged bags securely.
- 6 Store under dry conditions.

## Application rates

How to calculate the application rate with pots and containers >10 liter:

Total rate when using a 50-liter container:

3 g/l for first 10 liters = 30 gram

1 g/l for the additional 40 liters = 40 gram

Total = 30 + 40 = 70 gram per 50-liter container

	Containers < 10 liter	Containers > 10 liter
Container nursery stock	3-5 g/l	1g/l for every extra liter content

Attention Reduce these recommended rates for sensitive plants. Only use the product for topdressing. If you are using drip irrigation, place the product under the drippers. Irrigate well after application. This product is not suitable for mixing or dibbling in substrates. Topdress products cannot be used with capillary mats or ebb/flood irrigation systems. Contact your ICL advisor for more detailed advice. Trial first on a small scale before changing the rate, application or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results.

## Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.