



Osmocote® Start

Start

Optimum results for your salt-sensitive crops from start to finish

11 | 11 | 17 | 2.0 | TE
N P2O5 K2O MgO



Guaranteed analysis

Oxide		
N	Total Nitrogen	11%
	Nitrate nitrogen (N-NO3)	4.5%
	Ammoniacal nitrogen (N-NH4)	6.5%
P2O5	Phosphorus Pentoxide	11%
	Water soluble (P2O5)	8.2%
K2O	Potassium Oxide	17%
	Water Soluble (K2O)	17.0%
MgO	Magnesium Oxide	2.0%
	Water soluble (MgO)	1.0%
B	Boron	0.014%
	Water soluble (B)	0.011%
Cu	Copper	0.090%
	Water soluble (Cu)	0.080%
	Copper EDTA (Cu)	0.090%
Fe	Iron	0.38%
	Water soluble (Fe)	0.010%
	Iron EDTA (Fe)	0.15%
Mn	Manganese	0.055%
Mo	Molybdenum	0.013%
	Water soluble (Mo)	0.013%
Zn	Zinc	0.014%
	Water soluble (Zn)	0.014%

Description

Give your short growth cycle plants the best possible start with Osmocote® Start coated fertilizer, perfect for your violets, primulas, impatiens, marigolds, and many more. It's also ideal for your more salt-sensitive vegetable crops and for those cuttings proving difficult to root. As every single granule contains every element required, you run no risk of de-mixing any nutrients. The start of something great.

Benefits

- Improve crop color and root development
- Increased efficiency with reduced leaching
- Dust-free and easy to use

How to use

- 1 The longevity of Osmocote Start is determined at 21°C. At lower average soil temperatures, the product will work longer at higher average soil temperatures shorter. Indicational: 16°C: 5-7M, 21°C: 5-6M, 26°C: 4-5M.
- 2 You can mix Osmocote® Start and use it for topdressing, just make sure not to leave any granules stuck to the leaves.
- 3 Partly used bags must be closed / sealed properly.
- 4 Store under dry and cool conditions.
- 5 If you need more information, please contact your technical support.

Application rates

	Light feeding	Normal feeding
Container nursery stock	0.5-1.5 g/l	1.5-2.5 g/l
Pot plants	0.5-1.5 g/l	1.5-2.5 g/l
Bedding plants	0.5-2.0 g/l	-
Vegetable young plants	0.5-1.5 g/l	1.5-2.5 g/l

Attention These recommended rates are based on unfertilized substrates. Please be aware that these are general recommendations. Specific situations such as use in tunnels, greenhouses, or specific climate conditions require adjustments. This product is not recommended for dibbling. Contact your ICL Specialty Fertilizers advisor for more detailed advice. Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and the application of our products is beyond our control, ICL Specialty Fertilizers cannot be held responsible for any adverse results.

Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.