



Combifert®

18-10-12+2CaO+2MgO+18SO₃+0.2B+0.1Zn

A full package for all a plant could need

18 | 10 | 12 | 2 | 2 | 18 | TE
N P₂O₅ K₂O CaO MgO SO₃

Guaranteed analysis

Oxide		
N	Total Nitrogen	18%
	Ammoniacal nitrogen (N-NH ₄)	3.9%
	Urea nitrogen (N-Urea)	14.1%
P ₂ O ₅	Phosphorus Pentoxide	10%
	Water soluble (P ₂ O ₅)	9%
K ₂ O	Potassium Oxide	12%
	Water Soluble (K ₂ O)	12%
CaO	Calcium Oxide	2%
	Water soluble (CaO)	2%
MgO	Magnesium Oxide	2%
	Water soluble (MgO)	2.0%
SO ₃	Sulphur trioxide	18%
	Water soluble (SO ₃)	17%
B	Boron	0.2%
	Water soluble (B)	0.2%
Zn	Zinc	0.1%
	Water soluble (Zn)	0.1%

Description

Combifert® 18-10-12+2CaO+2MgO+18SO₃+B+Zn is chock-full of goodness. There's the balanced blend of NPK you'd expect, but that's not all. A high content of sulfur for improved uptake of vitamins and proteins, calcium for stronger cells and roots, magnesium to aid photosynthesis, plus boron for improved cell wall formation and sugar movement, and finally zinc for chlorophyll production. It's the perfect mix for your crops – for plant health, good development, and quality fruit and vegetables.

Benefits

- \\ Balanced NPK plus sulfur, calcium, magnesium, boron, and zinc
- \\ Easy to apply
- \\ Tailor-made for more efficient nutrition

How to use

- 1 Apply evenly to the surface of the soil, close to the root system, and then scratch into the soil.
- 2 Alternatively, localize the application and bury the granules in rows close to the root system.
- 3 Store under dry conditions.
- 4 Properly seal partly used or damaged bags.
- 5 For more information or recommendations, please contact your nearest ICL distributor or the ICL advisor for your area.

Application rates

The average recommended application rate is 300 to 700 kg of fertilizer per hectare, depending on the type of crop and expected yield.

Trail first on a small scale before changing the rate, or any other variables, As circumstances can differ and the application of our products is beyond our control, ICL cannot be held responsible for any adverse results.

Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.
