



Combifert®

Citrus

Bring a zing to your citrus grove

20 | 10 | 5 | 2.0 | 13.0 | TE
N P2O5 K2O MgO SO3



Guaranteed analysis

Oxide		
N	Total Nitrogen	20%
	Ammoniacal nitrogen (N-NH4)	3.5%
	Urea nitrogen (N-Urea)	16.5%
P2O5	Phosphorus Pentoxide	10%
	Water soluble (P2O5)	6.5%
K2O	Potassium Oxide	5%
	Water Soluble (K2O)	5.0%
MgO	Magnesium Oxide	2.0%
	Water soluble (MgO)	2.0%
SO3	Sulphur trioxide	13.0%
	Water soluble (SO3)	13.0%
Fe	Iron	3.30%

Description

Combifert® Citrus is a special formula developed for citrus trees, which can need more iron and magnesium than other crops. It contains 3.3% iron and 2% magnesium, as well as a citrus-friendly blend of NPK. It's also rich in sulfur (13%) which trees need for amino acid synthesis and chlorophyll formation, and can also lower your soil's pH level, which means improved uptake of phosphor and microelements for your trees. This granular fertilizer is proven to be effective for all your citrus needs: beautiful, juicy fruit will be all yours.

Benefits

- \\ Specially formulated for citrus trees
- \\ Easy to apply
- \\ Iron, magnesium, and sulfur to suit citrus tree requirements

How to use

- 1 Apply evenly to the surface of the soil, close to the root system, and then scratch into the soil. Alternatively, localize the application and bury the granules in rows close to the root system.
- 2 We recommend applying Combifert® Citrus between rows at least 15 to 20 days before bud-break to ensure the tree can access all the necessary nutrients during this period of high nutritional demand.
- 3 Store under dry conditions.
- 4 Properly seal partly used or damaged bags.
- 5 For more information or recommendations, please contact your nearest ICL distributor or the ICL advisor for your area.

Application rates

Trail first on a small scale before changing the rate, or any other variables, As circumstances can differ and the application of our products is beyond our control, ICL cannot be held responsible for any adverse results.

Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.