



Agrocote[®]
Max

Max 43-0-0

Hit the nitro for Max-imum speedy growth

43 | 0 | 0
N P₂O₅ K₂O



Guaranteed analysis

Oxide		
N	Total Nitrogen	43%
	Urea nitrogen (N-Urea)	43.0%
P ₂ O ₅	Phosphorus Pentoxide	0%
K ₂ O	Potassium Oxide	0%

Description

Give your crops that nitrogen kick and boost growth with Agrocote[®] Max 43-0-0 | 3-4M. Ideal for pre-planting or pre-seeding application in light, Mediterranean-area soil conditions, this high-nitrogen formulation is perfectly suited to your short-medium cycle crops, particularly open field-grown vegetables such as potatoes. Thanks to ICL's state of the art patented E-Max Release Technology, you can rest assured that your plants will enjoy a controlled high-quality, urea-based nitrogen release over a guaranteed 3-4 month period.

Benefits

- Increases crop yield or maintains it in case of lower nutrient application rate
- Enhanced nitrogen use efficiency
- Simple all-in-one base fertilizer solution

How to use

- 1 Apply by row-application or broadcasting.
- 2 Do not apply more than 1-2 weeks prior to planting.
- 3 Incorporate into the soil at a depth of between 5-10 cm.
- 4 Place the product under drippers where applicable.
- 5 Improve product and plant performance by irrigating after planting/seeding.
- 6 Reduce the dosage if WSFs are used in the second part of the crop cycle.
- 7 Properly seal partly used or damaged bags.
- 8 Store under dry conditions.

Application rates

Crops:	Kg/ha
Potato (processing and consumption):	150-400 kg/ha
Fruits and open field vegetables:	150-400 kg/ha

Suitable for short-medium crop cycles in Mediterranean areas. Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and the application of our products is beyond our control, ICL cannot be held responsible for any adverse results.

Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.