



**Agrocote®
Max**

Max 44-0-0

Put the N in uNrivalled growth

44 | 0 | 0
N P2O5 K2O



Guaranteed analysis

Oxide		
N	Total Nitrogen	44%
	Urea nitrogen (N-Urea)	44.0%
P2O5	Phosphorus Pentoxide	0%
K2O	Potassium Oxide	0%

Description

Give your crops all the nitrogen they could ever possibly need with Agrocote® Max 44-0-0 | 2-3M. One single application in lighter soil conditions, prior to planting or seeding, is enough to effortlessly fulfil the substantial nitrogen requirements of your potatoes and any other medium-late cycle field soil-grown vegetables. With ICL's state of the art patented E-Max Release Technology, your crops will benefit from a controlled, reliable, and sustained high-quality nitrogen release over a pre-determined 2-3 month period, based on an average soil temperature of 21°C.

Benefits

- \\ One single base fertilizer instead of multiple applications
- \\ Increases crop yield or maintains it in case of lower nutrient application rate
- \\ Enhanced nitrogen use efficiency

How to use

- 1 Apply by row application or broadcasting.
- 2 Do not apply more than 1-2 weeks prior to planting.
- 3 Incorporate into the soil at a depth of between 5-10 cm.
- 4 Place the product under drippers where applicable.
- 5 Improve product and plant performance by irrigating after planting/seeding.
- 6 Reduce the dosage if WSFs are used in the second part of the crop cycle.
- 7 Properly seal partly used or damaged bags.
- 8 Store under dry conditions.
- 9 If you need more information, please contact your technical support.

Application rates

Crops:	Kg/ha
Potato (processing and consumption):	150-400 kg/ha
Fruits and open field vegetables:	150-400 kg/ha

Suitable for medium-late crop cycles. Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and the application of our products is beyond our control, ICL cannot be held responsible for any adverse results.

Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.