

Solinure® FX

8-8-38+2MgO+TE

Feel the FX of perfect fresh fruit

8 | 8 | 38 | 2,0 | TE N P205 K20 Mg0





Guaranteed analysis

Total Nitrogen	8%
Ammoniacal nitrogen (N-NH4)	1.1%
Urea nitrogen (N-Urea)	6.9%
Phosphorus Pentoxide	8%
Water soluble (P2O5)	8.0%
Potassium Oxide	38%
Water Soluble (K2O)	38.0%
Magnesium Oxide	2.0%
Water soluble (MgO)	2.0%
Boron	0.01%
Water soluble (B)	0.01%
Copper	0.002%
Water soluble (Cu)	0.002%
Copper EDTA (Cu)	0.002%
Iron	0.04%
Water soluble (Fe)	0.04%
Iron EDTA (Fe)	0.04%
Manganese	0.01%
Water soluble (Mn)	0.01%
Manganese EDTA (Mn)	0.01%
Molybdenum	0.002%
Water soluble (Mo)	0.002%
Zinc	0.002%
Water soluble (Zn)	0.002%
Zinc EDTA (Zn)	0.002%
	Urea nitrogen (N-Urea) Phosphorus Pentoxide Water soluble (P2O5) Potassium Oxide Water Soluble (K2O) Magnesium Oxide Water soluble (MgO) Boron Water soluble (B) Copper Water soluble (Cu) Copper EDTA (Cu) Iron Water soluble (Fe) Iron EDTA (Fe) Manganese Water soluble (Mn) Manganese EDTA (Mn) Molybdenum Water soluble (Mo) Zinc Water soluble (Zn)

Description

Drip irrigation is a useful practice to enhance water and nutrient use efficiency. The use of drip irrigation significantly reduces nitrous oxide (N2O) emissions compared to furrow and sprinkler irrigation systems.

Benefits

- Dependability
- High K formula with magnesium and microelements
- Value for money: the FX line delivers all the necessary
- Clean materials: made with low impurity ingredients
- ICL quality: tight control of ingredients, manufacturing, and consistency



How to use

- Use Solinure® FX 30-10-10+2MgO+TE is especially recommended for the developing and production phases.
- 2 Store under dry conditions.
- 3 Make sure you properly seal partly used or damaged bags.
- For specific advice and recommendations, contact ICL or your professional advisor.

Application rates

Recommended rate: Apply 12-22kg/1000 m2 per week.

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and the application of our products is beyond our control, ICL cannot be held responsible for any adverse results.

Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.

