



Organ 2-0-8

 A first-class eco-friendly solution

2

N

0

P₂O₅

8

K₂O



Description

Flecotec Organ 2-0-8 is a formulation with a high potassium ratio, containing 4% free amino acids obtained by enzymatic hydrolysis of vegetable proteins. Potassium plays an important role in fruit development, promoting quality, color, flavor and consistency. In the production of Flecotec Organ 2-0-8, high-quality plant material of non-GMO origin has been used, subjected to different fermentation processes in which the yeast *Corynebacterium Melassecola* intervenes.

Flecotec Organ 2-0-8 can be used in organic farming in accordance with Regulation (EU) 2018/848 of the European Parliament and of the Council.

Certificates for organic farming



Guaranteed analysis

Oxide

N	Total Nitrogen	2%
	Ammoniacal nitrogen (N-NH ₄)	0.7%
	Organic nitrogen	1.3%
 P ₂ O ₅	Phosphorus Pentoxide	0%
K ₂ O	Potassium Oxide	8%
	Water Soluble (K ₂ O)	8%

Benefits

- Rich in potassium
- Incorporates amino acids
- It can be used in organic farming in accordance with Regulation (EU) 2018/848

Characteristics

EC at 1 g/l:	0.35 mS/cm
pH:	5.5-6.5
Density (g/cc):	1.16
Container:	20 L carafe, 1000 L container and bulk tank

Application rates

You can apply Flecotec Organ 2-0-8 to most crops and soils by fertigation with a rate of 15-25 l/ha per application at times of peak potassium demand.

These recommended dosages are provided as a guideline only, as growing conditions may vary depending on the species grown, varieties, irrigation water, soil, climate, and overall crop health. Please contact your ICL delegate if you need further advice on dosage.

Do a first trial or small-scale trial before modifying the dose, application, or any other variables. As circumstances may vary and we have no control over the application of our products, ICL cannot be held liable in the event of adverse results.



How to use

- 1 The use of Flecotec Organ 2-0-8 is recommended in the stages of maximum potassium demand
- 2 Store in a dry place at temperatures between 5 and 30°C.
- 3 If you need more information, ask for technical advice.

Recommended period of use

J	F	M	A	M	J	J	A	S	O	N	D

Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.

<https://icl-growingsolutions.com/agriculture/products/flecotec-organ-2-0-8/>
: 12/05/2024