



Agrolution[®] Liquid

ME-5

The liquid solution for ME-5



Guaranteed analysis

Oxide		
N	Total Nitrogen	0%
P2O5	Phosphorus Pentoxide	0%
K2O	Potassium Oxide	0%
Cu	Copper	0.150%
	Water soluble (Cu)	0.150%
	Copper EDTA (Cu)	0.150%
Fe	Iron	3.65%
	Water soluble (Fe)	3.65%
	Iron EDTA (Fe)	3.65%
Mn	Manganese	1.70%
	Water soluble (Mn)	1.70%
	Manganese EDTA (Mn)	1.70%
Mo	Molybdenum	0.140%
	Water soluble (Mo)	0.140%
Zn	Zinc	0.340%
	Water soluble (Zn)	0.340%
	Zinc EDTA (Zn)	0.340%

Characteristics

Description

Cater to your horticultural, fruit, and extensive crops' every micro-element need with Agrolution[®] Liquid ME-5. While rich in iron, manganese, copper, and zinc chelated with EDTA, along with mineral-form molybdenum, this boron-free fertigation fertilizer is perfect for your irrigation systems comprising a 0.3 ppm B content and above. Use this versatile solution as a complement to any simple fertilizer base or to bolster any micro element-free soluble NPK fertilizer applications.

Benefits

- \\ Prevents and corrects plant nutritional deficiencies
- \\ Designed for fertigation fertilizer application
- \\ Made from high-quality formulations

How to use

- 1 Do not mix Agrolution Liquid ME-5 with oil-based products or others capable of significantly altering the solution's pH level.
- 2 We recommend first conducting a small-scale trial or consulting your representative, ICL technical advisor, or distributor.

Application rates

Crops and stages	Crop Treatment period	No. of applications
Fruit trees, grapevines, olive trees, citrus fruit trees	From early bud-break to the fruit setting stage	3-6 fertigations at a rate of 20-30 l/Ha per year
Horticultural crops	Apply first treatment when leaves cover at least 2/3 of the cultivated area	Apply through fertigation at rate of 25-40 l/Ha per year

These recommended rates are only a guide, as crop conditions can vary depending on cultivated species, variety, irrigation water, soil, climate, and overall crop health. Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and the application of our products is beyond our control, ICL cannot be held responsible for any adverse results.

Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.