



# Combifert®

12-12-17+3CaO+3MgO+15SO<sub>3</sub>

A beautiful balance: NPK, sulfur, calcium, and magnesium

12 | 12 | 17 | 3 | 3 | 15  
N P<sub>2</sub>O<sub>5</sub> K<sub>2</sub>O CaO MgO SO<sub>3</sub>

## Guaranteed analysis

Oxide		
N	Total Nitrogen	12%
	Ammoniacal nitrogen (N-NH <sub>4</sub> )	3.4%
	Urea nitrogen (N-Urea)	8.6%
P <sub>2</sub> O <sub>5</sub>	Phosphorus Pentoxide	12%
	Water soluble (P <sub>2</sub> O <sub>5</sub> )	9%
K <sub>2</sub> O	Potassium Oxide	17%
	Water Soluble (K <sub>2</sub> O)	17%
CaO	Calcium Oxide	3%
	Water soluble (CaO)	3%
MgO	Magnesium Oxide	3%
	Water soluble (MgO)	3.0%
SO <sub>3</sub>	Sulphur trioxide	15%
	Water soluble (SO <sub>3</sub> )	12%

## Description

Combifert® 12-12-17+3CaO+3MgO+15SO<sub>3</sub> will give your fruit trees, vegetables, or arable crops a direct blast of balanced nutrients. There's NPK plus a high sulfur content for improved uptake, as well as calcium for stronger cells and roots, and magnesium for more efficient photosynthesis. It's a simple-to-use, homogeneous fertilizer for good healthy development.

## Benefits

- \\ A balanced blend without too much nitrogen
- \\ Easy to apply
- \\ Tailor-made for more efficient nutrition

## How to use

- 1 Apply evenly to the surface of the soil, close to the root system.
- 2 Scratch into the soil.
- 3 Alternatively, localize the application and bury the granules in rows close to the root system.
- 4 Apply Combifert 12-12-17+3CaO+3MgO+15SO<sub>3</sub> 10 to 20 days before early bud-break in tree crops and 7 to 10 days before sowing or planting arable and horticultural crops.
- 5 For more information or recommendations, please contact your nearest ICL distributor or the ICL advisor for your area.

## Application rates

---

300 to 700 kg of fertilizer per hectare, depending on the type of crop and expected yield.

Trail first on a small scale before changing the rate, or any other variables, As circumstances can differ and the application of our products is beyond our control, ICL cannot be held responsible for any adverse results.

## Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.