



Mg-7

Give your crops the magnesium and sulfur they need for healthy growth

0 | 0 | 0 | 7 | 14
N P2O5 K2O MgO SO3



Guaranteed analysis

Oxide		
N	Total Nitrogen	0%
P2O5	Phosphorus Pentoxide	0%
K2O	Potassium Oxide	0%
MgO	Magnesium Oxide	7%
	Water soluble (MgO)	7.0%
SO3	Sulphur trioxide	14%
	Water soluble (SO3)	14%

Description

Treat your crops to an effective magnesium-rich boost with Flecotec® Mg-7 fertilizer. This specially designed formula is the perfect solution for remedying any magnesium or sulfur deficiencies in any of your crops. By increasing available levels of magnesium, you will be ensuring enhanced photosynthesis and carbohydrate creation in your plants. Providing additional sulfur, on the other hand, will boost your plants' vital processes. Flecotec Mg-7 can be used in ecological agriculture in accordance with Council Regulation (CE) 2018/848. In European countries, consult the applicable local regulations for using Flecotec Mg-7 as a certified ecological product, or contact your ICL adviser in the region.

Benefits

- Rich in magnesium for enhanced photosynthesis
- Rich in sulfur for enhance amino-acid synthesis
- Can be used in ecological agriculture in accordance with Council Regulation (CE) 2018/848

How to use

- 1 Use Flecotec® Mg-7 to prevent and remedy sulfur and magnesium deficiencies in all types of crops. Only use when necessary.
- 2 Do not exceed the recommended dose.
- 3 While you can mix this product with most commonly-used fertilizers and plant protection products, you should always carry out a compatibility test prior to application.
- 4 Do not apply this product in high temperatures.
- 5 Do not mix with oils or calcium-based products.
- 6 Store under dry conditions, between 5-30 °C.
- 7 If you need more information, please contact your technical support.

Recommended period of use

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NNOV	DEC

Application rates

Recommended rates

Organic farming	Only apply Flecotec® Mg-7 when you are certain that the prevention or correction of magnesium deficiencies is required, applying at a rate of 20-30 l/ha per application.
Intensive crops	Apply between 150-250 l/ha per crop cycle. Spread out across several applications according to the crops' specific needs.
Foliar spray	To remedy magnesium deficiencies (magnesium chlorosis), apply 5-10 cm³/l of water (0.5-1 l/hl) and re-apply every ten days.

Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.