



Magnesium

Put the Mag in magnificent growth

0 | 0 | 0 | 16 | 32
N P2O5 K2O MgO SO3



Guaranteed analysis

Oxide		
N	Total Nitrogen	0%
P2O5	Phosphorus Pentoxide	0%
K2O	Potassium Oxide	0%
MgO	Magnesium Oxide	16%
	Water soluble (MgO)	16.0%
SO3	Sulphur trioxide	32%
	Water soluble (SO3)	32%

Characteristics

Description

Give your fruit, horticultural, and extensive crops all the magnesium they are crying out for with Flecotec® Magnesium fertilizer. Your crops will enjoy enhanced photosynthesis thanks to this highly-soluble product's high magnesium content. With the addition of boosted levels of sulfur, you can optimize each and every plant's vital processes, and therefore guarantee all-round healthy growth. Flecotec Magnesium can be used in ecological agriculture in accordance with Council Regulation (CE) 2018/848. In European countries, consult the applicable local regulations for using Flecotec Magnesium as a certified ecological product, or contact your ICL adviser in the region.

Benefits

- \\ Efficient source of sulfur and magnesium
- \\ Nitrogen free for a rebalanced fertigation program
- \\ Can be used in ecological agriculture in accordance with Council Regulation (CE) 2018/848

How to use

- 1 Use Flecotec® Magnesium to prevent and correct sulfur and magnesium deficiencies in all crop types. Only use when necessary.
- 2 Do not exceed the recommended dose.
- 3 Store under dry conditions, between 5-30 °C.
- 4 Properly seal partly used or damaged bags.
- 5 If you need more information, please contact us for technical support.

Recommended period of use

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC

Application rates

Recommended rates

Organic farming	Only apply Flecotec® Magnesium when you are certain that the prevention or correction of magnesium deficiencies is required, applying at a rate of 25-30 kg/ha per application.
Intensive crops	Apply 25-30 kg/ha per application. Apply 4-5 times throughout the crop cycle.
Extensive crops	Apply between 120-130 kg/ha per crop cycle, spread out across several applications.
Foliar spray	To remedy magnesium deficiencies (magnesium chlorosis), apply 0.2-0.3 kg/hl and re-apply every ten days.

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and the application of our products is beyond our control, ICL cannot be held responsible for any

Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.