



# Combifert®

6-9-22+2MgO+21SO<sub>3</sub>

A winning combination for effective growth

6 | 9 | 22 | 2 | 21  
N P<sub>2</sub>O<sub>5</sub> K<sub>2</sub>O MgO SO<sub>3</sub>

## Guaranteed analysis

Oxide		
N	Total Nitrogen	6%
	Ammoniacal nitrogen (N-NH <sub>4</sub> )	6%
P <sub>2</sub> O <sub>5</sub>	Phosphorus Pentoxide	9%
	Water soluble (P <sub>2</sub> O <sub>5</sub> )	8.5%
K <sub>2</sub> O	Potassium Oxide	22%
	Water Soluble (K <sub>2</sub> O)	22%
MgO	Magnesium Oxide	2%
	Water soluble (MgO)	2.0%
SO <sub>3</sub>	Sulphur trioxide	21%
	Water soluble (SO <sub>3</sub> )	21%

## Description

Combifert® 6-9-22+2MgO+21SO<sub>3</sub> contains a mix of NPK that is extra rich in potassium: excellent for bringing your crops to perfect fruition and improving water efficiency. It also packs a big punch of sulfur for enhanced protein and vitamin processing in your plants, plus it contains magnesium for better photosynthesis. Easy to use with vegetables, fruits, or extensive arable crops for healthier, happier plants.

## Benefits

- High potassium and sulfur content, plus magnesium
- Easy to apply
- Tailor-made for more efficient nutrition

## How to use

- 1 Apply evenly to the surface of the soil, close to the root system, and then scratch into the soil.
- 2 Alternatively, localize the application and bury the granules in rows close to the root system.
- 3 Store under dry conditions.
- 4 Properly seal partly used or damaged bags.
- 5 For more information or recommendations, please contact your nearest ICL distributor or the ICL advisor for your area.

## Application rates

---

The average recommended application rate is 300 to 700 kg of fertilizer per hectare, depending on the type of crop and expected yield.

Trail first on a small scale before changing the rate, or any other variables, As circumstances can differ and the application of our products is beyond our control, ICL cannot be held responsible for any adverse results.

## Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.