



# Agroblen®

18-5-11+2MgO

Give your crops the very best

18 | 5 | 11 | 2.0  
N P2O5 K2O MgO



## Guaranteed analysis

Oxide		
N	Total Nitrogen	18%
	Nitrate nitrogen (N-NO <sub>3</sub> )	7.5%
	Ammoniacal nitrogen (N-NH <sub>4</sub> )	8.4%
	Urea nitrogen (N-Urea)	2.1%
P <sub>2</sub> O <sub>5</sub>	Phosphorus Pentoxide	5%
	Water soluble (P <sub>2</sub> O <sub>5</sub> )	3.7%
K <sub>2</sub> O	Potassium Oxide	11%
	Water Soluble (K <sub>2</sub> O)	11.0%
MgO	Magnesium Oxide	2.0%

## Description

Reap the fruits of your labor with Agroblen® 18-5-11+2MgO | 3-4M, the controlled release fertilizer specifically designed for your strawberry and soft fruit crops' nutritional needs. This product's high-nitrogen fully coated NPK formula with added magnesium ensures fast growth and efficient crop establishment, even in wider fruit and vegetable crops. Following just one simple row application, you can sit back and enjoy 3-4 months of controlled nutrition for your plants thanks to ICL's patented Resin Release Technology.

## Benefits

- High in nitrogen for uniform growth
- Reduced nutrient leaching during irrigation or rainfall
- Designed specifically for strawberry production established by "frigo" plantlets

## How to use

- 1 Apply by row application or broadcasting.
- 2 Do not apply more than 1-2 weeks before planting.
- 3 Incorporate into the soil at a depth between 5-10 cm.
- 4 Place the product under drippers where applicable.
- 5 Improve product and plant performance by irrigating after application.
- 6 Reduce the applied dosage if WSFs are used in the second part of the crop cycle.
- 7 Store under dry conditions.
- 8 Properly seal partly used or damaged bags.
- 9 If you need more information, please contact your technical support.

## Application rates

<b>Crops:</b>	<b>Rate</b>
Strawberries:	12-15 g/plant for open field soil grown crops (at 30,000 pl/ha)
Strawberries:	3-4 g/plant for open field soil grown crops, in beds (at 100,000 pl/ha)
Other soft fruits:	300-450 kg/ha, row application
Other suitable crops:	300-600 kg/ha

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and the application of our products is beyond our control, ICL cannot be held responsible for any adverse results.

## Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.