



# Agromaster®

22-10-10+MgS

Steady release for even growth

22 | 10 | 10 | 4.0 | 10.0  
N P2O5 K2O MgO SO3



## Guaranteed analysis

| Oxide |                             |       |
|-------|-----------------------------|-------|
| N     | Total Nitrogen              | 22%   |
|       | Ammoniacal nitrogen (N-NH4) | 3.0%  |
|       | Urea nitrogen (N-Urea)      | 19.0% |
| P2O5  | Phosphorus Pentoxide        | 10%   |
|       | Water soluble (P2O5)        | 7.5%  |
| K2O   | Potassium Oxide             | 10%   |
|       | Water Soluble (K2O)         | 10.0% |
| MgO   | Magnesium Oxide             | 4.0%  |
| SO3   | Sulphur trioxide            | 10.0% |

## Description

Agromaster® 22-10-10+4MgO+10SO<sub>3</sub> | 5-6M is formulated with an NPK ratio of 2-1-1, plus plenty of magnesium and sulfur, so your crops will enjoy consistent, uniform growth and better photosynthesis. It's ideal for medium to long crop cycles as it contains extremely high levels of coated NPK. Our combined E-Max and Resin Release Technologies ensure nutrients are available when they're needed, over 5 – 6 months, whatever the weather conditions might throw at you and particularly in spring when fertilizing is often difficult. This formula is especially suitable for vegetables in protected areas, such as cucumbers, tomatoes, eggplants, or peppers, or in Mediterranean or subtropical areas. Agromaster® releases nutrients more efficiently than conventional fertilizers so more gets to your plants: any nutrient loss is minimized so your yields will be more fruitful.

## Benefits

- Controlled release NPK for months of uniform growth, plus magnesium and sulfur
- Higher yield, or use less and still maintain your usual yield
- Minimizes nutrient losses, especially in light soils

## How to use

- 1 Apply in spots, beds, or row application.
- 2 Incorporate Agromaster® in the soil between 5-10cm depth.
- 3 Where applicable, place the product under drippers.
- 4 Irrigating after application improves product and plant performance.
- 5 If you're using WSFs in the second part of the crop cycle, reduce the dosage of Agromaster® accordingly.
- 6 Store under dry conditions.
- 7 Properly seal partly used or damaged bags.
- 8 If you need more information, please contact your technical support.

## Application rates

| Crops   | Rates         |
|---|---------------|
| Medium-long crop cycle vegetables grown in protected areas: Tomato, pepper, eggplant, cucumber, and gherkin | 400-800 kg/ha |
| Fruits and vegetables grown in open field in Mediterranean and sub-tropical areas                           | 400-800 kg/ha |

Trial first on a small scale before changing the rate, or any other variables. As circumstances can differ and the application of our products is beyond our control, ICL cannot be held responsible for any adverse results.

## Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.