



Combifert®

18-6-6+6CaO+3.6MgO+25SO₃

Full speed ahead for healthy growth

18 | 6 | 6 | 6 | 3.6 | 25
N P₂O₅ K₂O CaO MgO SO₃

Guaranteed analysis

Oxide		
N	Total Nitrogen	18%
	Ammoniacal nitrogen (N-NH ₄)	2.3%
	Urea nitrogen (N-Urea)	15.7%
P ₂ O ₅	Phosphorus Pentoxide	6%
	Water soluble (P ₂ O ₅)	5.6%
K ₂ O	Potassium Oxide	6%
	Water Soluble (K ₂ O)	6%
CaO	Calcium Oxide	6%
	Water soluble (CaO)	6%
MgO	Magnesium Oxide	3.6%
	Water soluble (MgO)	3.6%
SO ₃	Sulphur trioxide	25%
	Water soluble (SO ₃)	25%

Description

Combifert® 18-6-6+6CaO+3.6MgO+25SO₃ is a great blend of everything your crops could need at their peak growing stage. It's a carefully selected blend of NPK that's super-rich in nitrogen, with sulfur for improved nutrient uptake, calcium for stronger cells and roots, and magnesium for improved photosynthesis. A tailor-made compound fertilizer ideal for your fruit, vegetable, or arable crops.

Benefits

- Rich in nitrogen and sulfur, plus PK, calcium, and magnesium
- Easy to apply
- Tailor-made for more efficient nutrition

How to use

- 1 Apply evenly to the surface of the soil, close to the root system, and then scratch into the soil.
- 2 Alternatively, localize the application and bury the granules in rows close to the root system.
- 3 Store under dry conditions.
- 4 Properly seal partly used or damaged bags.
- 5 For more information or recommendations, please contact your nearest ICL distributor or the ICL advisor for your area.

Application rates

The average recommended application rate is 300 to 700 kg of fertilizer per hectare, depending on the type of crop and expected yield.

Trail first on a small scale before changing the rate, or any other variables, As circumstances can differ and the application of our products is beyond our control, ICL cannot be held responsible for any adverse results.

Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.
