



Topdress FT 4-5M

Top-of-the-line solution for all your topdress application needs

22 | 5 | 6 | 2.0 | TE
N P₂O₅ K₂O MgO



Guaranteed analysis

Oxide		
N	Total Nitrogen	22%
	Nitrate nitrogen (N-NO ₃)	4.8%
	Ammoniacal nitrogen (N-NH ₄)	5.9%
	Urea nitrogen (N-Urea)	11.3%
P ₂ O ₅	Phosphorus Pentoxide	5%
	Water soluble (P ₂ O ₅)	3.7%
K ₂ O	Potassium Oxide	6%
	Water Soluble (K ₂ O)	6.0%
MgO	Magnesium Oxide	2.0%
	Water soluble (MgO)	1.0%
Cu	Copper	0.050%
Fe	Iron	0.80%
Mn	Manganese	0.30%
Zn	Zinc	0.100%

Description

Osmocote® Topdress FT 4-5M has been designed with your container nursery stock in mind. It's made to meet all your topdress application needs. It incorporates special Fusion Technology to ensure it adheres easily to your growing medium. So if your containers blow over, no granules will be lost.

Benefits

- \\ High in fast- and slow-release nitrogen and phosphorous
- \\ Ensures even distribution with fine granule size
- \\ Contains added trace elements for optimal greening effect

How to use

- 1 Reduce these recommended rates for sensitive plant material.
- 2 Apply the product only by topdressing. In case drip irrigation is used, place the product under the drippers.
- 3 This product is not meant for mixing or dibbling in substrates.
- 4 Topdress products cannot be used with capillary mats or ebb / flow irrigation systems.
- 5 Store under dry conditions.
- 6 Properly seal partly used or damaged bags.

Application rates

How to calculate the application rate with pots and containers >10 liter:

Total rate when using a 50 liter container:

3 g/l for first 10 liter = 30 gram

1 g/l for the additional 40 liter = 40 gram

Total = 30 + 40 = 70 gram per 50 liter container

	Containers < 10 liters	Containers > 10 liters
Container Nursery Stock	2 – 4 g/l	for every extra liter content

Trial first on a small scale before changing the rate, or any other variables. As circumstances can differ and the application of our products is beyond our control, ICL cannot be held responsible for any adverse results.

Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.