

Preseeder

Perfect preseeding fertilizer

18 | 22,0 | 5, N P205 K20









Guaranteed analysis

| Oxio | le | |
|------|-----------------------------|-------|
| N | Total Nitrogen | 18% |
| | Ammoniacal nitrogen (N-NH4) | 6.6% |
| | Urea nitrogen (N-Urea) | 2.8% |
| | Ureaformaldehyde (N-MU) | 8.6% |
| P20 | 5 Phosphorus Pentoxide | 22.0% |
| | Water soluble (P2O5) | 18.7% |
| K20 | Potassium Oxide | 5.% |

Description

When you're overseeding or laying new turf, there's no better solution than Sierraform® GT Preseeder. It contains both slow-release nitrogen and slow-release potassium, along with conventional release NPK (18-22-5) in a single compound micro-granule. The high phosphorus analysis in Preseeder promotes good rooting and establishment on your greens, tees, fairways, and sports fields, while minimizing salt levels to boost germination rates. Its nutrient release pattern is tailored to the needs of new turf or seed.

Benefits

- High P analysis to support seeding, overseeding and sod laying operations
- Slow release N (MU) for consistent and sustained turf response
- **1** Even spread at low application rates



How to use

- Use from March to September.
- Apply to dry foliage and irrigate after 1 2 days if there is no rainfall. Irrigation helps with dispersion and lowers risk of mower pick-up on close cut surfaces. Avoid applying during frosty or drought conditions.
- Delay any vertical cutting and/or grooming until 3 days after application, to allow the granules to properly disperse. Significantly less nitrogen is released when soil temperatures are below 10° C.
- If you need more information, please contact your technical support.

Recommended period of use

| JAN | FEB | MAR | APR | MAY | JUN | JUL | AUG | SEP | OCT | NNOV | DEC |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|------|-----|
| | | | | | | | | | | | |

Application rates

Recommended Rate:

15-30g/m2

Trial first on a small scale before changing the rate, or any other variables. As circumstances can differ and the application of our products is beyond our control, ICL cannot be held responsible for any adverse results.

Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.

