

Ficote[®] Total

8-9M

Your go-to for a long-term base dressing

17 | 9 | 11 | 2,0 | TE N P205 K20 MgO



Guaranteed analysis

Oxide	•	
N	Total Nitrogen	17%
	Nitrate nitrogen (N-NO3)	7.6%
	Ammoniacal nitrogen (N-NH4)	9.4%
P2O5	Phosphorus Pentoxide	9%
	Water soluble (P2O5)	6.6%
K20	Potassium Oxide	11%
	Water Soluble (K2O)	11.0%
MgO	Magnesium Oxide	2.0%
	Water soluble (MgO)	1.3%
В	Boron	0.013%
	Water soluble (B)	0.011%
Cu	Copper	0.032%
	Water soluble (Cu)	0.020%
Fe	Iron	0.25%
	Iron EDTA (Fe)	0.05%
Mn	Manganese	0.03%
	Water soluble (Mn)	0.035%
Мо	Molybdenum	0.012%
	Water soluble (Mo)	0.012%
Zn	Zinc	0.015%
	Water soluble (Zn)	0.010%

Description

Longevity is not a problem with Ficote® Total 8-9M as it's a 2nd-generation coated fertilizer. Ficote® Total is designed to be used as a basal feed, and you can combine it with a water-soluble fertilizer. Contains all the trace elements your plants will need as well as 100% coated NPK: it's also great value. If your release pattern during the longevity is crucial, we recommend Osmocote® Exact patterned release fertilizer. This range is the epitome of safety and uniformity in release.

Benefits

- Nerfect for base dressings
- Nutrients when plants need it, and great value for money
- Coated NPK plus all essential trace elements



How to use

- Ficote® Total is designed for base dressing and can be combined with a water-soluble fertilizer.
- We recommend using Osmocote® Exact for difficult growing conditions, growing crops in covered areas, salt sensitive crops, high value crops, plant hole dibbling and when using full rates of coated fertilizer.
- The longevity of Ficote Total is determined at 21°C. At lower average soil temperatures, the product will work longer at higher average soil temperatures shorter. Indicational: 16°C: 5-7M, 21°C: 5-6M, 26°C: 4-5M.
- Partly used bags must be closed / sealed properly.
- 5 Store under dry and cool conditions.
- 6 If you need more information, please contact your technical support.

Application rates

Container nursery stock

Light feeding: 2-2.5 g/l

Normal feeding: 3-3.5 g/l

Heavy feeding: 4-4.5 g/l

Attention The recommended rates mentioned are based on unfertilized substrates. Please note that these are general recommendations. Contact your ICL Specialty Fertilizers advisor for more detailed advice. Trial first on a small scale before changing the rate, application or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL Specialty Fertilizers cannot be held responsible for any adverse results.

Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.

