

Sierrablen Plus

Active 3M

Active growth for all activity

19 | 5 | 18 | 2,0 | TE N P205 K20 Mg0








Guaranteed analysis

| Oxide | | |
|---------------------------|-----------------------------|--------|
| Ν | Total Nitrogen | 19% |
| | Ammoniacal nitrogen (N-NH4) | 1.2% |
| | Urea nitrogen (N-Urea) | 15.6% |
| P2O5 Phosphorus Pentoxide | | 5% |
| | Water soluble (P2O5) | 3.7% |
| K20 | Potassium Oxide | 18% |
| | Water Soluble (K2O) | 18.0% |
| MgO | Magnesium Oxide | 2.0% |
| Cu | Copper | 0.010% |
| Mn | Manganese | 0.10% |
| Zn | Zinc | 0.030% |

Description

When seeking active growth for active endeavors, look no further than Sierrablen® Plus Active 3M. This high nitrogen and potassium controlled release fertilizer is ideal for spring and fall applications. With ICL's patented Poly-S coating technology, your turf will benefit from a controlled 3-month nutrient release. Aid stress recovery and improve overall plant health thanks to its full trace element package. Boost your turf's color and hide stress symptoms thanks to its added magnesium content.

Benefits



Improves turf color and overall health

Masks stress symptoms and aids recovery



How to use

- Apply directly to dry foliage.
- Watering-in after application helps dispersion and boosts immediate effect. Do not apply during drought or frosty conditions.
- You should apply following completion of aeration programs, including scarification, slitting and/or hollow tining to prevent damage to the granules.
- 4 Properly seal partly used or damaged bags.
- 5 Store under dry conditions.
- 6 If you need more information, please contact your technical support.

Recommended period of use

JAN FEB MAR APR MAY JUN JUL AUG SEP OCT NNOV DEC

Application rates

Recommended Rate:

25.00 - 35.00 g/m²

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and the application of our products is beyond our control, ICL cannot be held responsible for any adverse results.

Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.

