



Sierraform[®] GT

CalMag

For soil low in calcium and magnesium

14 | 0 | 7 | 8 | 5 | TE
N P2O5 K2O CaO MgO



Guaranteed analysis

Oxide		
N	Total Nitrogen	14%
	Ammoniacal nitrogen (N-NH4)	1.3%
	Urea nitrogen (N-Urea)	2.9%
	Ureaformaldehyde (N-MU)	9.8%
P2O5	Phosphorus Pentoxide	0%
K2O	Potassium Oxide	7%
	Water Soluble (K2O)	7%
CaO	Calcium Oxide	8%
	Water soluble (CaO)	8%
MgO	Magnesium Oxide	5%
	Water soluble (MgO)	5.0%
Cu	Copper	0.02%
Fe	Iron	0.5%
Mn	Manganese	0.100%
Mo	Molybdenum	0.001%
Zn	Zinc	0.02%

Description

An agronomically balanced N to K ratio, with additional calcium and magnesium. Containing trace elements and SMX for improved stress tolerance and stronger roots, and sustained-release MU-based nitrogen for regular, steady growth. You'll get even coverage with these compound mini granules, even applied at a low rate. Ideal throughout the summer on all fine turf. Healthier grass all round, with nutrients released for up to 8 – 10 weeks.

Benefits

- 2:1 N:K ratio to support active in-season growth
- Ca & Mg supplied for all round turf health
- Stronger roots, healthier grass, steady growth
- Slow release N (MU) for consistent and sustained turf response

How to use

- 1 Apply to dry foliage and irrigate after 1 – 2 days if there is no rainfall.
- 2 Irrigation helps with dispersion and lowers risk of mower pick-up on close cut surfaces.
- 3 Watering-in after application will minimize footprint trails. Avoid applying during frosty or drought conditions.
- 4 Delay any vertical cutting and/or grooming until 3 days after application, to allow the granules to properly disperse. Significantly less nitrogen is released when soil temperatures are below 10° C.
- 5 If spilled on pavement, concrete, clothes, etc. brush off immediately as it can stain. Contains iron: do not apply in case of risk of contact with concrete, marble, stone, fabric, or swimming pools.
- 6 If you need more information, please contact your technical support.

Recommended period of use

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC

Application rates

Recommended Rate:

15 – 30 g/m²

Trial first on a small scale before changing the rate, or any other variables. As circumstances can differ and the application of our products is beyond our control, ICL cannot be held responsible for any adverse results.

Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.