



## Agroleaf® Special

### Zn

Do away with zinc deficiency thanks to this water-soluble fertilizer



## Guaranteed analysis

Oxide		
N	Total Nitrogen	0%
P <sub>2</sub> O <sub>5</sub>	Phosphorus Pentoxide	0%
K <sub>2</sub> O	Potassium Oxide	0%
Zn	Zinc	14.000%
	Water soluble (Zn)	14.000%
	Zinc EDTA (Zn)	14.000%

## Characteristics

## Description

Agroleaf® Special Zn is a premium foliar feed that prevents and controls zinc deficiency. It's suitable for a range of crops including cereals, vegetables, flowers, and fruit-trees. Plus, it's fully water-soluble, making it a popular choice in agriculture and horticulture. It contains a specific amino acid known as X<sub>3</sub> Active, which aids the crop in absorbing Special Zn quickly. This reduces physiological stress caused by zinc deficiency within just two days. Preparing the solution is simple, and it fully dissolves in no time. We use only the finest raw materials and latest technologies to produce Agroleaf® Special Zn, which contains the highest concentration of active compound in its class. You can also tank-mix Special Zn with other fertilizers and crop-protection products, saving you time out on the field.

## Benefits

- \\ Correct Zn deficiency
- \\ Absorbed quickly and effectively to combat stress
- \\ Boosts plant metabolism

## How to use

- 1 The best time for spraying is late afternoon or evening. Avoid spraying in unfavorable conditions (for example water-stressed crops, low relative air humidity, high temperatures, or if rainfall is expected within 3 days). If rainfall occurs within 4 hours of spraying, re-spray crop 3 or 4 days later.
- 2 Test the product in its final concentration on a small plot of several plants, one week before the actual commercial spraying. Check results several days later.
- 3 Do the same especially when intending to tank-mix with other substances.
- 4 Use a high-pressure sprayer equipped with nozzles that produce very fine droplets.
- 5 Fill the sprayer tank to half its capacity; start the agitation mechanism. Put the entire amount of intended Agroleaf® Special Zn in the tank and wait until it completely dissolves. Put the full intended amount of fungicides and/or insecticides in the tank. Make sure that the pH is between 4 and 6.5; correct if necessary by adding very small amounts of Nova® PeKacid.
- 6 Fill the sprayer tank up to its full capacity. Start spraying immediately and don't stop until the tank is empty. Do not store spraying solution overnight.
- 7 If you need more information, please contact your technical support.

## Application rates

---

The average recommended application rate is 0.2 to 0.3% dosage for all crops

Trial first on a small scale before changing the rate, or any other variables, As circumstances can differ and the application of our products is beyond our control, ICL cannot be held responsible for any adverse results.

## Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.

---