

Agrolution[®] Special

212

Give your plants Special treatment

23	10	23 TE
Ν	P2O5	K2O



Guaranteed analysis

Oxide		
N	Total Nitrogen	23%
	Nitrate nitrogen (N-NO3)	5.2%
	Urea nitrogen (N-Urea)	17.8%
P2O5	Phosphorus Pentoxide	10%
	Water soluble (P2O5)	10.0%
К2О	Potassium Oxide	23%
	Water Soluble (K2O)	23.0%
В	Boron	0.01%
	Water soluble (B)	0.01%
Cu	Copper	0.010%
	Water soluble (Cu)	0.010%
	Copper EDTA (Cu)	0.010%
Fe	Iron	0.16%
	Water soluble (Fe)	0.16%
	Iron DTPA (Fe)	0.16%
Mn	Manganese	0.08%
	Water soluble (Mn)	0.08%
	Manganese EDTA (Mn)	0.08%
Мо	Molybdenum	0.006%
	Water soluble (Mo)	0.006%
Zn	Zinc	0.040%
	Water soluble (Zn)	0.040%
	Zinc EDTA (Zn)	0.040%

Characteristics

Description

For those crops that command a balanced nitrogen and phosphorus-rich NPK-based nutrition, choose Agrolution® Special 212 fertigation fertilizer. With this tailored application, your plants can enjoy all the nutrients they could ever need, with their nutrient availability also improving after each use. Thanks to its pure composition, high solubility, and acidifying effect, you can prevent any possible blockages and deposits within your irrigation system. This product is also formulated to remedy any calcium deficiencies within your crops.

Benefits

- Completely soluble
- Enhances nutrient availability
- Nemedies nitrogen, potassium, and calcium deficiencies





How to use



3

The Agrolution® range offers a wide range of analyses covering all crop growth stages while providing tailored assistance to specific soil and water types.

2 Store under dry conditions.

Properly seal partly used or damaged bags.

4 For tailored advice, contact your ICL advisor and ask for information specific to Agrolution® Special.

Application rates

The average recommended application rate is 40-50 kg/ha per week.

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and the application of our products is beyond our control, ICL cannot be held responsible for any adverse results.

Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.

