



Solinure® GT

7

Help your plants get going

18 | 11 | 11 | 2.0 | TE
N P2O5 K2O MgO



Guaranteed analysis

Oxide		
N	Total Nitrogen	18%
	Nitrate nitrogen (N-NO3)	3.3%
	Ammoniacal nitrogen (N-NH4)	11.3%
	Urea nitrogen (N-Urea)	3.4%
P2O5	Phosphorus Pentoxide	11%
	Water soluble (P2O5)	11.0%
K2O	Potassium Oxide	11%
	Water Soluble (K2O)	11.0%
MgO	Magnesium Oxide	2.0%
	Water soluble (MgO)	2.0%
B	Boron	0.01%
	Water soluble (B)	0.01%
Cu	Copper	0.002%
	Water soluble (Cu)	0.002%
	Copper EDTA (Cu)	0.002%
Fe	Iron	0.04%
	Water soluble (Fe)	0.04%
	Iron EDTA (Fe)	0.04%
Mn	Manganese	0.01%
	Water soluble (Mn)	0.01%
	Manganese EDTA (Mn)	0.01%
Mo	Molybdenum	0.002%
	Water soluble (Mo)	0.002%
Zn	Zinc	0.002%
	Water soluble (Zn)	0.002%
	Zinc EDTA (Zn)	0.002%

Description

Give your greenhouse and tunnel system crops their ideal streamlined nutrient kick with Solinure® GT 7 fertigation fertilizer. This high-nitrogen NPK formula has been specifically designed with your greenhouse and tunnel system plants in mind, particularly tailored for application during their vegetative growth stage. With minimal urea and no chloride present, your plants will enjoy a fully chelated trace element package, along with added magnesium for all-round healthy growth.

Benefits

- \\ Ideal for greenhouse and tunnel system application
- \\ Designed for plants' vegetative growth stage
- \\ Highly dependable product

Characteristics

How to use

- 1 Use Solinure® GT 7 for vegetative stages of crop growth.
- 2 Store under dry conditions.
- 3 Properly seal partly used or damaged bags.
- 4 For specific advice and recommendations, contact ICL or your professional advisor.

Application rates

Crops:	Kg/ha
Open field and protected area crops and vegetables:	40-60 kg/ha per week
Berry crops, trees, soft and stone fruits:	30-50 kg/ha per week

Recommended rate: Apply 4-5 kg/1000m² per week Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and the application of our products is beyond our control, ICL cannot be held responsible for any adverse results.

Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.