



Solinure® GT

5

Helps your crops get through the winter with flying colors

20 | 20 | 20 | TE
N P2O5 K2O

5

Guaranteed analysis

Oxide		
N	Total Nitrogen	20%
	Nitrate nitrogen (N-NO ₃)	5.9%
	Ammoniacal nitrogen (N-NH ₄)	3.8%
	Urea nitrogen (N-Urea)	10.3%
P2O5	Phosphorus Pentoxide	20%
	Water soluble (P2O5)	20.0%
K2O	Potassium Oxide	20%
	Water Soluble (K2O)	20.0%
B	Boron	0.01%
	Water soluble (B)	0.01%
Cu	Copper	0.002%
	Water soluble (Cu)	0.002%
	Copper EDTA (Cu)	0.002%
Fe	Iron	0.04%
	Water soluble (Fe)	0.04%
	Iron EDTA (Fe)	0.04%
Mn	Manganese	0.01%
	Water soluble (Mn)	0.01%
	Manganese EDTA (Mn)	0.01%
Mo	Molybdenum	0.002%
	Water soluble (Mo)	0.002%
Zn	Zinc	0.002%
	Water soluble (Zn)	0.002%
	Zinc EDTA (Zn)	0.002%

Characteristics

Description

For that all-round nutritional hit, look no further than Solinure® GT 5 fertigation fertilizer. With its balanced NPK formula, this ever-dependable fertilizer is your ideal solution for assisting your greenhouse and tunnel system plants in their demanding growing and ripening stages, while also being perfectly suited to helping your crops through those challenging winter production periods. With as little urea as possible and completely chloride-free, your plants are sure to benefit from its fully cheated trace element package and added magnesium.

Benefits

- \\ Balanced NPK formula
- \\ Highly dependable
- \\ Designed for growing and ripening stage application

How to use

- 1 Use Solinure® GT 5 for growing and ripening stages.
- 2 Store under dry conditions.
- 3 Properly seal partly used or damaged bags.
- 4 For specific advice and recommendations, contact ICL or your professional advisor.

Application rates

Crops:	Kg/ha
Open field and protected area crops and vegetables:	40-60 kg/ha per week
Berry crops, trees, soft and stone fruits:	30-50 kg/ha per week

Recommended rate: Apply 4-5 kg/1000m² per week Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and the application of our products is beyond our control, ICL cannot be held responsible for any adverse results.

Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.