



Solinure® GT

4

Give your plants the nutrition they need, when they need it

14 | 6 | 23 | 2.0 | TE
N P2O5 K2O MgO

4

Guaranteed analysis

Oxide		
N	Total Nitrogen	14%
	Nitrate nitrogen (N-NO3)	6.1%
	Ammoniacal nitrogen (N-NH4)	7.9%
P2O5	Phosphorus Pentoxide	6%
	Water soluble (P2O5)	6.0%
K2O	Potassium Oxide	23%
	Water Soluble (K2O)	23.0%
MgO	Magnesium Oxide	2.0%
	Water soluble (MgO)	2.0%
B	Boron	0.01%
	Water soluble (B)	0.01%
Cu	Copper	0.002%
	Water soluble (Cu)	0.002%
	Copper EDTA (Cu)	0.002%
Fe	Iron	0.04%
	Water soluble (Fe)	0.04%
	Iron EDTA (Fe)	0.04%
Mn	Manganese	0.01%
	Water soluble (Mn)	0.01%
	Manganese EDTA (Mn)	0.01%
Mo	Molybdenum	0.002%
	Water soluble (Mo)	0.002%
Zn	Zinc	0.002%
	Water soluble (Zn)	0.002%
	Zinc EDTA (Zn)	0.002%

Description

When your fruit and vegetable crops are crying out for their favorite nutrient fix, look no further than Solinure® GT 4 fertigation fertilizer. Complete with added trace element package, this high-potassium, low-phosphorus NPK formula is designed to help your greenhouse and tunnel system-grown plants through their growing and ripening stages, even during those cold and dark winter months. Give your plants only what they actually need, with a solution that is both free from chlorides and contains minimal urea.

Benefits

- Designed for greenhouse and tunnel system application
- High-potassium, low-phosphorus NPK ratio
- Completely chloride-free

Characteristics

How to use

- 1 Use Solinure® GT 4 for growing and ripening stages, along with winter crop production.
- 2 Store under dry conditions.
- 3 Properly seal partly used or damaged bags.
- 4 For specific advice and recommendations, contact ICL or your professional advisor.

Application rates

Crops:	Kg/ha
Open field and protected area crops and vegetables:	40-60 kg/ha per week
Berry crops, trees, soft and stone fruits:	30-50 kg/ha per week

Recommended rate: Apply 4-5 kg/1000m² per week Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and the application of our products is beyond our control, ICL cannot be held responsible for any adverse results.

Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.