



Solinure® GT

8

Give your crops the very best

23 | 10 | 10 | 5.6 | TE
N P2O5 K2O MgO

8

Guaranteed analysis

Oxide		
N	Total Nitrogen	23%
	Nitrate nitrogen (N-NO3)	1.0%
	Ammoniacal nitrogen (N-NH4)	1.2%
	Urea nitrogen (N-Urea)	20.8%
P2O5	Phosphorus Pentoxide	10%
	Water soluble (P2O5)	10.0%
K2O	Potassium Oxide	10%
	Water Soluble (K2O)	10.0%
MgO	Magnesium Oxide	5.6%
	Water soluble (MgO)	5.6%
B	Boron	0.01%
	Water soluble (B)	0.01%
Cu	Copper	0.002%
	Water soluble (Cu)	0.002%
	Copper EDTA (Cu)	0.002%
Fe	Iron	0.04%
	Water soluble (Fe)	0.04%
	Iron EDTA (Fe)	0.04%
Mn	Manganese	0.01%
	Water soluble (Mn)	0.01%
	Manganese EDTA (Mn)	0.01%
Mo	Molybdenum	0.002%
	Water soluble (Mo)	0.002%
Zn	Zinc	0.002%
	Water soluble (Zn)	0.002%
	Zinc EDTA (Zn)	0.002%

Description

Don't compromise when it comes to your crops' nitrogen-thirsty nutritional needs, simply choose Solinure® GT 8 fertigation fertilizer. With its high-nitrogen NPK ratio, this highly dependable formula is your perfect solution for the vegetative growing stages of your greenhouse and tunnel system crops. This chloride-free product contains added magnesium, along with a full trace element package, providing your plants with an optimized formula that promotes all-round healthy growth.

Benefits

- \\ Completely chloride free
- \\ Helps plants through vegetative stages of crop growth
- \\ Complete with full trace element package

Characteristics

How to use

- 1 Use Solinure® GT 8 for the vegetative stages of plant growth.
- 2 Store under dry conditions.
- 3 Properly seal partly used or damaged bags.
- 4 For specific advice and recommendations, contact ICL or your professional advisor.

Application rates

Crops:	Kg/ha
Open field and protected area crops and vegetables:	40-60 kg/ha per week
Berry crops, trees, soft and stone fruits:	30-50 kg/ha per week

Recommended rate: Apply 4-5 kg/1000m² per week Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and the application of our products is beyond our control, ICL cannot be held responsible for any adverse results.

Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.