



Solinure® GT

2

Keep your fruit crops flourishing during every stage and season

7 | 19 | 38 | 2.0 | TE
N P₂O₅ K₂O MgO

2

Guaranteed analysis

Oxide		
N	Total Nitrogen	7%
	Nitrate nitrogen (N-NO ₃)	7.0%
P ₂ O ₅	Phosphorus Pentoxide	19%
	Water soluble (P ₂ O ₅)	19.0%
K ₂ O	Potassium Oxide	38%
	Water Soluble (K ₂ O)	38.0%
MgO	Magnesium Oxide	2.0%
	Water soluble (MgO)	2.0%
B	Boron	0.01%
	Water soluble (B)	0.01%
Cu	Copper	0.002%
	Water soluble (Cu)	0.002%
	Copper EDTA (Cu)	0.002%
Fe	Iron	0.04%
	Water soluble (Fe)	0.04%
	Iron EDTA (Fe)	0.04%
Mn	Manganese	0.01%
	Water soluble (Mn)	0.01%
	Manganese EDTA (Mn)	0.01%
Mo	Molybdenum	0.002%
	Water soluble (Mo)	0.002%
Zn	Zinc	0.002%
	Water soluble (Zn)	0.002%
	Zinc EDTA (Zn)	0.002%

Characteristics

Description

Solinure® GT 2 fertigation fertilizer is a high-potassium, low-nitrogen NPK formula. It's your ideal solution for helping your greenhouse and tunnel system-grown fruit crops not only through their growing and ripening stages, but also through those cool, dark winter months. While free from chlorides and urea, your plants will be boosted by this dependable product's added magnesium, along with its fully chelated trace element package.

Benefits

- \\ Perfect for fruit growing and ripening stages
- \\ Designed for greenhouse and tunnel system application
- \\ Includes added magnesium and trace elements

How to use

- 1 Use Solinure® GT 2 for the growing and ripening stages of fruit, along with winter crop production.
- 2 Store under dry conditions.
- 3 Properly seal partly used or damaged bags.
- 4 For specific advice and recommendations, contact ICL or your professional advisor.

Application rates

Crops:	kg/ha
Open field and protected area crops and vegetables:	40-60 kg/ha per week
Berry crops, trees, soft and stone fruits:	30-50 kg/ha per week

Recommended rate: Apply 4-5 kg/1000m² per week Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and the application of our products is beyond our control, ICL cannot be held responsible for any adverse results.

Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.