



Solinure® GT

3

Get the most out of your fruit crops with Solinure® GT 3

12 | 5 | 35 | 2.0 | TE
N P2O5 K2O MgO

3

Guaranteed analysis

Oxide		
N	Total Nitrogen	12%
	Nitrate nitrogen (N-NO3)	8.1%
	Ammoniacal nitrogen (N-NH4)	2.1%
	Urea nitrogen (N-Urea)	1.8%
P2O5	Phosphorus Pentoxide	5%
	Water soluble (P2O5)	5.0%
K2O	Potassium Oxide	35%
	Water Soluble (K2O)	35.0%
MgO	Magnesium Oxide	2.0%
	Water soluble (MgO)	2.0%
B	Boron	0.01%
	Water soluble (B)	0.01%
Cu	Copper	0.002%
	Water soluble (Cu)	0.002%
	Copper EDTA (Cu)	0.002%
Fe	Iron	0.04%
	Water soluble (Fe)	0.04%
	Iron EDTA (Fe)	0.04%
Mn	Manganese	0.01%
	Water soluble (Mn)	0.01%
	Manganese EDTA (Mn)	0.01%
Mo	Molybdenum	0.002%
	Water soluble (Mo)	0.002%
Zn	Zinc	0.002%
	Water soluble (Zn)	0.002%
	Zinc EDTA (Zn)	0.002%

Characteristics

Description

Get the most out of your tunnel system and greenhouse-grown fruit crops with Solinure® GT 3 fertigation fertilizer. With its low-nitrogen, high-potassium NPK formula, this is your perfect solution when it comes to your fruits' growing and ripening stages, along with boosting them through periods of winter production. Containing added magnesium, along with a fully chelated trace element package, this highly dependable product is free from unwanted chlorides and urea, providing your plants with only what they actually need.

Benefits

- Helps through dark, cool winter months
- Free from urea and chlorides
- Includes magnesium and a full trace element package

How to use

- 1 Use Solinure® GT 3 for the growing and ripening stages of fruit, along with winter crop production.
- 2 Store under dry conditions.
- 3 Properly seal partly used or damaged bags.
- 4 For specific advice and recommendations, contact ICL or your professional advisor.

Application rates

Crops:	Kg/ha
Open field and protected area crops and vegetables:	40-60 kg/ha per week
Berry crops, trees, soft and stone fruits:	30-50 kg/ha per week

Recommended rate: Apply 4-5 kg/1000m² per week Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results.

Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.