

Guaranteed analysis

Oxide

N	Total Nitrogen	20%
	Nitrate nitrogen (N-NO3)	5.6%
	Ammoniacal nitrogen (N-NH4)	3.9%
	Urea nitrogen (N-Urea)	10.5%
P2O5	Phosphorus Pentoxide	7%
	Water soluble (P2O5)	7.0%
K2O	Potassium Oxide	20%
	Water Soluble (K2O)	20.0%
MgO	Magnesium Oxide	2.0%
	Water soluble (MgO)	2.0%
В	Boron	0.01%
	Water soluble (B)	0.01%
Cu	Copper	0.002%
	Water soluble (Cu)	0.002%
	Copper EDTA (Cu)	0.002%
Fe	Iron	0.04%
	Water soluble (Fe)	0.04%
	Iron EDTA (Fe)	0.04%
Мо	Molybdenum	0.002%
	Water soluble (Mo)	0.002%
Zn	Zinc	0.002%
	Water soluble (Zn)	0.002%
	Zinc EDTA (Zn)	0.002%

Characteristics

Solinure[®] GT

30

The solution for all-round healthy plant growth

20 7 20 2.0 TE N P205 K20 Mg0

Description

When your greenhouse and tunnel system-grown fruit and vegetable crops find themselves in a bit of a rut, look no further than Solinure® GT 30 fertigation fertilizer. With its low-phosphorus NPK formula, along with added trace elements and magnesium, this cost-effective fertilizer is your ideal tool to help your plants through their starter stages and encourage better root development. Free from unwanted urea and chlorides, this consistently dependable solution contains added magnesium and trace elements for all-round healthy plant growth.

Benefits

- Consistently reliable product
- N High-quality materials with minimal impurities
- Fantastic value for money



How to use

4



- 2 Store under dry conditions.
- 3 Make sure you properly seal partly used or damaged bags.
 - For specific advice and recommendations, contact ICL or your professional advisor.

Application rates

Crops:	Kg/ha
Open field and protected area crops and vegetables:	40-60 kg/ha per week
Berry crops, trees, soft and stone fruits:	30-50 kg/ha per week

Recommended rate: Apply 3-10 kg/1000m² per week Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results.

Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.

