



Solinure® GT

31

Show your crop what you've GT

17 | 8 | 27 | 3.0 | TE
N P2O5 K2O CaO

Guaranteed analysis

Oxide		
N	Total Nitrogen	17%
	Nitrate nitrogen (N-NO3)	8.9%
	Ammoniacal nitrogen (N-NH4)	8.1%
P2O5	Phosphorus Pentoxide	8%
	Water soluble (P2O5)	8.0%
K2O	Potassium Oxide	27%
	Water Soluble (K2O)	27.0%
CaO	Calcium Oxide	3.0%
	Water soluble (CaO)	3.0%
B	Boron	0.01%
	Water soluble (B)	0.01%
Cu	Copper	0.002%
	Water soluble (Cu)	0.002%
	Copper EDTA (Cu)	0.002%
Fe	Iron	0.04%
	Water soluble (Fe)	0.04%
	Iron EDTA (Fe)	0.04%
Mn	Manganese	0.01%
	Water soluble (Mn)	0.01%
	Manganese EDTA (Mn)	0.01%
Mo	Molybdenum	0.002%
	Water soluble (Mo)	0.002%
Zn	Zinc	0.002%
	Water soluble (Zn)	0.002%
	Zinc EDTA (Zn)	0.002%

Description

For that calcium-rich nutrient boost your tunnel system and greenhouse-grown crops crave, look no further than Solinure® GT 31 fertigation fertilizer. With its high-potassium, low-phosphorus NPK ratio, see to any potential calcium deficiencies in your crops thanks to added calcium, complemented by a full trace element package. This fine balance is also designed to help your plants successfully complete their reproductive stages. Free from unwanted chlorides and urea, this cost-effective solution provides only what your plants actually need.

Benefits

- \\ Highly reliable product
- \\ Smart design provides exact nutrients required
- \\ Added calcium and trace elements for complete nutrition

Characteristics

How to use

- 1 Use Solinure® GT 31 for the reproductive stage, especially in the instance of calcium deficiencies.
- 2 Store under dry conditions.
- 3 Properly seal partly used or damaged bags.
- 4 For specific advice and recommendations, contact ICL or your professional advisor.

Application rates

Crops:	Kg/ha
Open field and protected area crops and vegetables:	40-60 kg/ha per week
Berry crops, trees, soft and stone fruits:	30-50 kg/ha per week

Recommended rate: Apply 4-10 kg/1000m² per week Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and the application of our products is beyond our control, ICL cannot be held responsible for any adverse results.

Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.