



# Solinure® FX

17

Feel the powerful FX of a dependable fertilizer

15 | 5 | 30  
N | P<sub>2</sub>O<sub>5</sub> | K<sub>2</sub>O



## Guaranteed analysis

Oxide		
N	Total Nitrogen	15%
	Ammoniacal nitrogen (N-NH <sub>4</sub> )	4.9%
	Urea nitrogen (N-Urea)	10.1%
P <sub>2</sub> O <sub>5</sub>	Phosphorus Pentoxide	5%
	Water soluble (P <sub>2</sub> O <sub>5</sub> )	5.0%
K <sub>2</sub> O	Potassium Oxide	30%
	Water Soluble (K <sub>2</sub> O)	30.0%

## Characteristics

## Description

When your open-field and protected area fruit and vegetable crops are crying out for a dependable, streamlined nutrient kick, simply reach for Solinure® FX 17 fertigation fertilizer. Although high in urea, this high-potassium NPK formula is chloride-free and contains no added trace elements. Providing excellent value for money, this acidifying Solinure product has the added benefit of ensuring that your drip lines are kept clean following application.

## Benefits

- \\ Contains urea, but is chloride-free
- \\ Ideal for open field soil-based application
- \\ No added trace elements

## How to use

- 1 Use Solinure® FX 17 for the growing and ripening stages of fruit.
- 2 Store under dry conditions.
- 3 Properly seal partly used or damaged bags.
- 4 For specific advice and recommendations, contact ICL or your professional advisor.

## Application rates

Crops:	kg/ha
Open field and protected areas crops and vegetables:	40-60 kg/ha per week
Berry crops, trees, soft and stone fruits:	30-50 kg/ha per week

Average recommended rate: Apply 40-60 kg/10,000m<sup>2</sup> per week Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results.

## Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.