

# Agroblen®

17-8-9+3MqO

Palm oil growth in the palm of your hand

17 | 8 | 9 | 3,0 N P205 K20 Mg0







# **Guaranteed analysis**

Oxide		
N	Total Nitrogen	17%
	Nitrate nitrogen (N-NO3)	7.6%
	Ammoniacal nitrogen (N-NH4)	9.4%
P2O5	Phosphorus Pentoxide	8%
	Water soluble (P2O5)	6.0%
K20	Potassium Oxide	9%
	Water Soluble (K2O)	9.0%
MgO	Magnesium Oxide	3.0%

## **Description**

Keep healthy palm oil crop growth right in the palm of your hand with Agroblen® 17-8-9+3MgO | 16-18M. This high-nitrogen granular NPK fertilizer is designed to help you seamlessly establish uniformly-growing tropical crops, such as palm oil seedlings. With its added magnesium, your plants will enjoy optimized photosynthetic activity. Simply sit back and watch your plants develop over 16-18 months from a controlled, sustained, and high-quality nutrient release, all made possible thanks to ICL's patented Resin Release Technology.

## **Benefits**

- NPK ratio promotes uniform vegetative growth in longterm crop cycles
- Can be applied in planting holes just before planting
- ★ Controlled nutrient release for uniform crop growth
- Added magnesium for improved photosynthetic activity



#### How to use

- 1 Apply using planting hole method.
- 2 You can also use row, spots, or broadcasting application methods.
- 3 Do not apply more than 1-2 weeks before planting.
- Incorporate into the soil at a depth of between 5-10 cm.
- 5 Place the product under drippers where applicable.
- 6 Improve product and plant performance by irrigating after application.
- Store under dry conditions. Properly seal partly used or damaged bags.

# **Application rates**

Crop:	Rate
Oil Palm: Pre-nursery stage (split in two holes aside the seed):  4 g/	
Oil Palm: Main nursery:	50 g/plant
Oil Palm: First year planting:	300 g/plant

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and the application of our products is beyond our control, ICL cannot be held responsible for any adverse results.

### Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.

