

Agroblen®

17-9-8+4MgO

Spring your temperate or sub-tropical crops to life

7 9 8 4,0 N P205 K20 Mg0







Guaranteed analysis

Oxide		
N	Total Nitrogen	17%
	Nitrate nitrogen (N-NO3)	7.5%
	Ammoniacal nitrogen (N-NH4)	9.5%
P205	Phosphorus Pentoxide	9%
	Water soluble (P2O5)	6.7%
K20	Potassium Oxide	8%
	Water Soluble (K2O)	8.0%
MgO	Magnesium Oxide	4.0%

Description

When spring has sprung, make sure your temperate and sub-tropical crops follow suit with Agroblen® 17-9-8+4MgO | 8-9M. This high-nitrogen granular NPK fertilizer has been specially formulated to help you establish your new temperate and sub-tropical plantations, such as avocados and citrus. With added magnesium for enhanced photosynthesis, this product is ideal for spring applications. Thanks to ICL's patented Resin Release Technology, your plants can enjoy a controlled, sustained, and reliable 8-9 month nutrient release.

Benefits

Can be applied in planting holes just before planting

Uniform crop growth thanks to steady and sustained nutrient release

\(\) Extra magnesium for enhanced photosynthesis



How to use

- 1 Apply using planting hole method.
- 2 You can also use row, spots, or broadcasting application methods.
- 3 Do not apply more than 1-2 weeks before planting.
- Incorporate into the soil at a depth of between 5-10 cm.
- 5 Place the product under drippers where applicable.
- 6 Improve product and plant performance by irrigating after application.
- Reduce the applied dosage if WSFs are used in the second part of the crop cycle.
- 8 Store under dry conditions. Properly seal partly used or damaged bags.

Application rates

Crops:	Rate
Orchards (new plantations):	75-100 g/plant
Avocado: plants grown in 3-liter pots. Application method: backfill (mixed with soil around throots):	ne 50 g/plant
Avocado: plants grown in 6-liter pots. Application method: backfill (mixed with soil around throots):	ne 100 g/plant
Citrus (new plantations, with young plants from 4-6 liter polybag):	150-200 g/plant
Coffee (new plantations):	50-100 g/plant
Other suitable crops: Vineyards and soft fruits (blackberry, raspberry, currants) (new plantations):	10-20 g/plant

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and the application of our products is beyond our control, ICL cannot be held responsible for any adverse results.

Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.

