



**Agroleaf<sup>®</sup>  
Liquid**

## Man Z+

Manganese and zinc deficiencies remedied in one



## Guaranteed analysis

Oxide		
N	Total Nitrogen	0%
P2O5	Phosphorus Pentoxide	0%
K2O	Potassium Oxide	0%
Mn	Manganese	4.70%
	Water soluble (Mn)	4.70%
	Manganese EDTA (Mn)	4.70%
Zn	Zinc	1.300%
	Water soluble (Zn)	1.300%
	Zinc EDTA (Zn)	1.300%

## Characteristics

## Description

Two nutrient deficiencies corrected with one solution: Agroleaf<sup>®</sup> Liquid Man Z+. Contains manganese and zinc micronutrients in a chelated form, developed for application by foliar sprays and/or fertigation, for both arable and fruit tree crops. Man Z+ can be used to prevent or cure the manganese and zinc deficiency, with more manganese than its sister product Agroleaf<sup>®</sup> Liquid Zinc M+. Use as a foliar spray on arable crops as a preventative measure. It's best to apply Man Z+ during the period of greatest vegetative growth. In fruit trees, it is recommended during initiation and development of major breakouts, avoiding full bloom. When applied by fertigation, Man Z+ is recommended for use as a preventative application.

## Benefits

- \\ Two in one: corrects and prevents manganese and zinc deficiencies
- \\ More manganese than Agroleaf<sup>®</sup> Liquid Zinc M+
- \\ Suitable for both arable and fruit tree crops

## How to use

- 1 General: Perform 2-3 preventative treatments on arable crops during the period of greatest vegetative growth, and on woody crops coinciding with the start and development of major sprouting, avoiding full flowering periods.
- 2 Dilute at the rate of 200-350 cc/HI without exceeding 8 L/Ha per application. Apply maximum dose if deficiencies are observed.
- 3 Use a wetting agent.
- 4 Citrus foliar spray: For citrus crops in particular, apply an initial dose in spring once new shoots have appeared, when the sufficiently developed new leaves are 2/3 of their natural size. Apply again within 15 days, and then again during summer flowering.
- 5 For applications with a total use of less than 2000 L/Ha, use higher doses (350 cc/HI); higher volumes (up to a maximum of 3000 L/Ha), use a half dose (250 cc/HI).
- 6 Foliar spray on stone and pip fruit: Apply during periods of greatest vegetative growth, when the crop has sufficient leaf mass. The recommended dose is 150-250 cc/HI with a maximum use of 1000 L/Ha.
- 7 Vegetables: The recommended dose is 150-200 cc/HI with a maximum use of between 800 and 1000 L/Ha.
- 8 If you need more information, please contact your technical support.

## Application rates

Trail first on a small scale before changing the rate, or any other variables, As circumstances can differ and the application of our products is beyond our control, ICL cannot be held responsible for any adverse results.

## Attention

Trial first on a small scale before changing the rate, application, or any other variables. As circumstances can differ and as the application of our products is beyond our control, ICL cannot be held responsible for any adverse results. Contact your ICL advisor for more detailed advice.